

CHEN'S PRIMES AND TERNARY GOLDBACH PROBLEM

HONGZE LI AND HAO PAN

ABSTRACT. We prove that there exists a $k_0 > 0$ such that every sufficiently large odd integer n with $3 \mid n$ can be represented as $p_1 + p_2 + p_3$, where p_1, p_2 are Chen's primes and p_3 is a prime with $p_3 + 2$ has at most k_0 prime factors.

1. INTRODUCTION

Let $\mathcal{P}$ denote the set of all primes. Define

$$\mathcal{P}_k = \{n : n \in \mathbb{N} \text{ and } n \text{ has at most } k \text{ prime divisors}\}$$

and

$$\mathcal{P}_k^{(2)} = \{p \in \mathcal{P} : p + 2 \in \mathcal{P}_k\}.$$

The well-known twin primes conjecture asserts that $\mathcal{P}_1^{(2)}$ has infinitely many elements. Nowadays, the best result on the twin primes conjecture belongs to Chen [1], who proved that $\mathcal{P}_2^{(2)}$ has infinitely many elements. In fact, Chen proved that for sufficiently large x ,

$$|\{p \in \mathcal{P}_2^{(2)} : p \leq x, (p + 2, P(x^{1/10})) = 1\}| \gg \frac{x}{(\log x)^2},$$

where

$$P(z) = \prod_{p < z} p.$$

In Iwaniec's unpublished notes [10], the exponent $1/10$ can be improved to $3/11$. In [6], Green and Tao say a prime p is Chen's prime if $p \in \mathcal{P}_2^{(2)}$.

On the other hand, in 1937 Vinogradov [18] solved the ternary Goldbach problem and showed that every sufficiently large odd integer can be represented as the sum of three primes. Two years later, using Vinogradov's method, van der Corput [2] proved that the primes contain infinitely many non-trivial three term arithmetic progressions (3AP). In 1999, with the help of the vector sieve method, Tolev [15] proved that there exist infinitely many non-trivial 3APs $\{p_1, p_2, p_3\}$ of primes, satisfying $p_1 \in \mathcal{P}_4^{(2)}$, $p_2 \in \mathcal{P}_5^{(2)}$ and $p_3 \in \mathcal{P}_{11}^{(2)}$.

However, in [6], with the help of the Szemerédi theorem, their transference principle and a result of Goldston and Yıldırım, Green and Tao proved that the primes contain arbitrarily long non-trivial arithmetic progressions. Certainly this is a remarkable breakthrough in additive number theory. Furthermore, Green and Tao

2000 *Mathematics Subject Classification.* Primary 11P32; Secondary 11N36, 11P55.

Key words and phrases. Chen prime; Ternary Goldbach problem; Rosser's weight.

This work was supported by the National Natural Science Foundation of China (Grant No. 10771135).

also claimed that using their method, one can prove that Chen's primes contain arbitrarily long non-trivial arithmetic progressions. And for the 3APs of Chen's primes, they proposed a detail proof in [5].

Let us return to the ternary Goldbach problems. In [13], using Tolev's method, Peneva proved that every sufficiently large odd integer n with $3 \mid n$ can be represented as $n = p_1 + p_2 + p_3$ with $p_i \in \mathcal{P}_{k_i}^{(2)}$, $k_1 = k_2 = 5$ and $k_3 = 8$. Subsequently, Tolev [16] improved Peneva's result to $k_1 = 2$, $k_2 = 5$ and $k_3 = 7$. Recently, Meng [12] proved that every sufficiently large odd integer n with $3 \nmid n - 1$ can be represented as $n = p_1 + p_2 + p_3$, where p_1 is Chen's prime, $p_2 \in \mathcal{P}_3^{(2)}$ and $p_3 \in \mathcal{P}$ (not of special type!).

Of course, we wish to prove that every sufficiently large odd integer n with $3 \mid n$ can be represented as the sum of three Chen's primes. Unfortunately, as we shall see later, it seems not easy. The key of Green and Tao's proof in [5] is to transfer Chen's primes to a subset with positive density of $\mathbb{Z}_N = \mathbb{Z}/N\mathbb{Z}$ (where N is a large prime). But this density is too small. However, in the present paper, we shall prove the following result:

Theorem 1.1. *There exists a positive integer k_0 such that every sufficiently large odd integer n with $3 \mid n$ can be represented as $p_1 + p_2 + p_3$, where p_1, p_2 are Chen's primes and $p_3 \in \mathcal{P}_{k_0}^{(2)}$.*

In Sections 2 and 3, we shall estimate some exponential sums involving the primes of special type. The proof of Theorem 1.1 will be given in Section 4.

2. THE MINOR ARCS

Let $k_0 \geq 8$ be a fixed integer and $B = 6^9$. Suppose that n is a sufficiently large integer, $W > 0$ is an even integer with $W \leq (\log n)^B$, and $1 \leq b \leq W$ satisfies $(b(b+2), W) = 1$. Let $\mathbb{T}$ denote the torus $\mathbb{R}/\mathbb{Z}$. For $1 \leq q \leq (\log n)^B$, define

$$\mathfrak{M}_{a,q} = \{\alpha \in \mathbb{T} : |\alpha q - a| \leq (\log n)^B/n\}.$$

Let

$$\mathfrak{M} = \bigcup_{\substack{1 \leq a \leq q \leq (\log n)^B \\ (a,q)=1}} \mathfrak{M}_{a,q}$$

and $\mathfrak{m} = \mathbb{T} \setminus \mathfrak{M}$.

Let $D = n^{0.32}$ and $z_0 = n^{1/k_0}$. For square-free number $d = p_1 p_2 \cdots p_k$ with primes $p_1 > p_2 > \cdots > p_k$, define Rosser's weights with the order D by

$$\lambda_D^+(d) = \begin{cases} (-1)^k & \text{if } p_1 \cdots p_{2l} p_{2l+1}^3 < D \text{ for all } 0 \leq l \leq (k-1)/2, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$\lambda_D^-(d) = \begin{cases} (-1)^k & \text{if } p_1 \cdots p_{2l-1} p_{2l}^3 < D \text{ for all } 1 \leq l \leq k/2, \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to see that $\lambda_D^\pm(d) = 0$ if $d \geq D$. Let $F(s)$ and $f(s)$ denote the functions of linear sieve. The following lemma is a fundamental result in sieve method.

Lemma 2.1 (Iwaniec [8, 9]). *Suppose that $\mathcal{P}_*$ is any set of primes and ω is a multiplicative function satisfying:*

$$0 < \omega(p) < p \text{ for } p \in \mathcal{P}_*, \quad \omega(p) = 0 \text{ for } p \notin \mathcal{P}_*,$$

and

$$\prod_{z_1 \leq p < z_2} \left(1 - \frac{\omega(p)}{p}\right)^{-1} \leq \frac{\log z_2}{\log z_1} \left(1 + \frac{L}{\log z_1}\right)$$

for a constant $L > 0$ and for all $2 \leq z_1 \leq z_2$. Then we have

$$\begin{aligned} \prod_{p < z} \left(1 - \frac{\omega(p)}{p}\right) &\leq \sum_{d|P_*(z)} \lambda_D^+(d) \frac{\omega(d)}{d} \\ &\leq \prod_{p < z} \left(1 - \frac{\omega(p)}{p}\right) (F(s) + O(e^{\sqrt{L}-s} (\log D)^{-1/3})), \end{aligned} \quad (2.1)$$

provided that $2 \leq z \leq D$, where $s = \log D / \log z$ and

$$P_*(z) = \prod_{p \in \mathcal{P}_* \cap [1, z)} p.$$

Similarly,

$$\begin{aligned} \prod_{p < z} \left(1 - \frac{\omega(p)}{p}\right) &\geq \sum_{d|P_*(z)} \lambda_D^-(d) \frac{\omega(d)}{d} \\ &\geq \prod_{p < z} \left(1 - \frac{\omega(p)}{p}\right) (f(s) + O(e^{\sqrt{L}-s} (\log D)^{-1/3})), \end{aligned} \quad (2.2)$$

provided that $2 \leq z \leq \sqrt{D}$. Furthermore, for any square-free integer q ,

$$\sum_{d|q} \lambda_D^-(d) \leq \sum_{d|q} \mu(d) \leq \sum_{d|q} \lambda_D^+(d). \quad (2.3)$$

Define

$$S_{n,z_0}(\alpha) = \sum_{\substack{p \leq n \\ p \equiv b \pmod{W} \\ (p+2, P(z_0))=1}} e(\alpha(p-b)/W) \log p.$$

Clearly

$$S_{n,z_0}(\alpha) = \sum_{\substack{p \leq n \\ p \equiv b \pmod{W}}} e(\alpha(p-b)/W) \log p \sum_{d|(p+2, P(z_0))} \mu(d).$$

Let

$$S_{n,z_0}^\pm(\alpha) = \sum_{\substack{p \leq n \\ p \equiv b \pmod{W}}} e(\alpha(p-b)/W) \log p \sum_{d|(p+2, P(z_0))} \lambda_D^\pm(d).$$

Lemma 2.2. *For any $\alpha \in \mathbb{T}$, we have $|S_{n,z_0}^+(\alpha) - S_{n,z_0}(\alpha)| \leq S_{n,z_0}^+(0) - S_{n,z_0}(0)$ and $|S_{n,z_0}(\alpha) - S_{n,z_0}^-(\alpha)| \leq S_{n,z_0}(0) - S_{n,z_0}^-(0)$.*

Proof. By (2.3),

$$\begin{aligned} |S_{n,z_0}^+(\alpha) - S_{n,z_0}(\alpha)| &\leq \sum_{\substack{p \leq n \\ p \equiv b \pmod{W}}} \log p \left| \sum_{d|(p+2, P(z_0))} \lambda^+(d) - \sum_{d|(p+2, P(z_0))} \mu(d) \right| \\ &= S_{n,z_0}^+(0) - S_{n,z_0}(0). \end{aligned}$$

The proof of the second inequality is similar. $\square$

Let τ denote the divisor function. It is well-known

$$\sum_{d \leq X} \tau(d)^A \ll_A X (\log X)^{2^A-1}$$

and

$$\sum_{d \leq X} \frac{\tau(d)^A}{d} \ll_A (\log X)^{2^A}.$$

Lemma 2.3. *Suppose that $X, X', Y, Y', Z, Z' > 0$ satisfy $X \leq X' \leq 2X$, $Y \leq Y' \leq 2Y$ and $XY \leq Z \leq Z' \leq X'Y'$. For any $d \geq 1$, let u_d, v_d, w_d be complex numbers with $|u_d|, |v_d|, |w_d| \leq \tau(d)^A (\log(XY))^A$. Suppose that $1 \leq a \leq q$ with $(a, q) = 1$, and $\alpha \in \mathbb{T}$ with $|\alpha q - a| \leq 1/q$. Then*

$$\begin{aligned} &\sum_{X \leq x \leq X'} u_x \left| \sum_{\substack{Y \leq y \leq Y' \\ Z \leq xy \leq Z' \\ xy \equiv b \pmod{W}}} v_y e(\alpha(xy - b)/W) \sum_{\substack{d|xy+2 \\ d \leq D}} w_d \right| \\ &\ll_A XY (\log(DXYq))^{2^{2A+2}} \tau^{2A+2}(W) \left(\frac{1}{q} + \frac{D^2W}{X} + \frac{qW}{XY} \right)^{1/4} \\ &\quad + XY^{1/2} (\log(DXY))^{2^{2A+2}} \tau^{2A+1}(W) \end{aligned} \tag{2.4}$$

provided that $X \geq D^2W$. Furthermore, suppose that $0 \leq v_{y_1} \leq v_{y_2} \leq (\log(XY))^A$ for any y_1, y_2 with $y_1 \leq y_2$. Then

$$\begin{aligned} &\sum_{X \leq x \leq X'} u_x \left| \sum_{\substack{Y \leq y \leq Y' \\ Z \leq xy \leq Z' \\ xy \equiv b \pmod{W}}} v_y e(\alpha(xy - b)/W) \sum_{\substack{d|xy+2 \\ d \leq D}} w_d \right| \\ &\ll_A XY (\log(DXYq))^{6^{A+2}} \left(\frac{1}{q} + \frac{DW}{Y} + \frac{qW}{XY} \right)^{1/4} \end{aligned} \tag{2.5}$$

provided that $Y \geq DW$.

Proof. By the Cauchy-Schwarz inequality,

$$\begin{aligned}
& \left(\sum_{X \leq x \leq X'} u_x \left| \sum_{\substack{Y \leq y \leq Y' \\ Z \leq xy \leq Z' \\ xy \equiv b \pmod{W}}} v_y e(\alpha(xy - b)/W) \sum_{\substack{d|xy+2 \\ d \leq D}} w_d \right| \right)^2 \\
& \leq \left(\sum_{X \leq x \leq X'} |u_x|^2 \right) \left(\sum_{\substack{X \leq x \leq X' \\ d_1, d_2 \leq D}} w_{d_1} \overline{w_{d_2}} \sum_{\substack{Y \leq y_1, y_2 \leq Y' \\ Z \leq xy_1, xy_2 \leq Z' \\ xy_1, xy_2 \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i} \text{ for } i=1,2}} v_{y_1} \overline{v_{y_2}} e(\alpha x(y_1 - y_2)/W) \right) \\
& \ll_A X(\log X)^{6^A} \sum_{\substack{1 \leq b' \leq W, (b', W)=1 \\ Y \leq y_1, y_2 \leq Y' \\ y_1 \equiv y_2 \equiv b' \pmod{W} \\ d_1, d_2 \leq D}} v_{y_1} \overline{v_{y_2}} w_{d_1} \overline{w_{d_2}} \sum_{\substack{X \leq x \leq X' \\ Z \leq xy_1, xy_2 \leq Z' \\ xb' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i} \text{ for } i=1,2}} e(\alpha x(y_1 - y_2)/W).
\end{aligned}$$

We have

$$\begin{aligned}
& \sum_{\substack{X \leq x \leq X' \\ Z \leq xy_1, xy_2 \leq Z' \\ xb' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i} \text{ for } i=1,2}} e(\alpha x(y_1 - y_2)/W) \\
& = \sum_{\substack{\max\{X, Z/y_1, Z/y_2\} \leq x \leq \min\{X', Z'/y_1, Z'/y_2\} \\ xb' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i} \text{ for } i=1,2}} e(\alpha x(y_1 - y_2)/W) \\
& \ll \min \left\{ \frac{X}{[d_1, d_2, W]}, \frac{1}{\|\alpha[d_1, d_2, W](y_1 - y_2)/W\|} \right\},
\end{aligned}$$

where $\|\theta\| = \min\{|\theta - t| : t \in \mathbb{Z}\}$. Hence for each $1 \leq b' \leq W$ with $(b', W) = 1$, we have

$$\begin{aligned}
& \left| \sum_{\substack{Y \leq y_1, y_2 \leq Y' \\ y_1 \equiv y_2 \equiv b' \pmod{W} \\ d_1, d_2 \leq D}} v_{y_1} \overline{v_{y_2}} w_{d_1} \overline{w_{d_2}} \right| \left| \sum_{\substack{X \leq x \leq X' \\ Z \leq xy_1, xy_2 \leq Z' \\ xb' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i} \text{ for } i=1,2}} e(\alpha x(y_1 - y_2)/W) \right| \\
& \ll \sum_{\substack{1 \leq y_1, y_2 \leq Y' \\ y_1 \equiv y_2 \equiv b' \pmod{W} \\ d_1, d_2 \leq D}} |v_{y_1}| |v_{y_2}| |w_{d_1}| |w_{d_2}| \min \left\{ \frac{X}{[d_1, d_2, W]}, \frac{1}{\|\alpha[d_1, d_2, W](y_1 - y_2)/W\|} \right\} \\
& \ll Y(\log(XY))^{6^A} \left(\sum_{h \leq D^2 Y} \tau^{4A+3}(hW) \min \left\{ \frac{XY}{hW}, \frac{1}{\|\alpha h\|} \right\} + \frac{X}{W} \sum_{h \leq D^2} \frac{\tau^{4A+2}(hW)}{h} \right) \\
& \quad (\text{where } h = [d_1, d_2, W](y_1 - y_2)/W \text{ if } y_1 > y_2, \text{ and } h = [d_1, d_2, W]/W \text{ if } y_1 = y_2) \\
& \ll Y(\log(DXY))^{6^A+1} \tau^{4A+3}(W) \max_{H \leq D^2 Y} \sum_{H/2 \leq h \leq H} \tau^{4A+3}(h) \min \left\{ \frac{XY}{HW}, \frac{1}{\|\alpha h\|} \right\} \\
& \quad + \frac{XY \tau^{4A+2}(W)}{W} (\log(DXY))^{2^{4A+3}}.
\end{aligned}$$

Applying Lemma 2.2 of [17], for any $H \leq D^2 Y$,

$$\begin{aligned}
& \sum_{H/2 \leq h \leq H} \tau^{4A+3}(h) \min \left\{ \frac{XY}{HW}, \frac{1}{\|\alpha h\|} \right\} \\
& \leq \left(\sum_{h \leq H} \tau^{8A+6}(h) \right)^{1/2} \left(\frac{XY}{HW} \sum_{h \leq H} \min \left\{ \frac{XY}{hW}, \frac{1}{\|\alpha h\|} \right\} \right)^{1/2} \\
& \ll_A \left(\frac{X^2 Y^2 (\log(DYq))^{2^{8A+6}}}{W^2} \left(\frac{1}{q} + \frac{D^2 W}{X} + \frac{qW}{XY} \right) \right)^{1/2}.
\end{aligned}$$

This concludes the proof of (2.4).

Let us turn to (2.5). Clearly

$$\begin{aligned}
& \left(\sum_{X \leq x \leq X'} u_x \left| \sum_{\substack{Y \leq y \leq Y' \\ Z \leq xy \leq Z' \\ xy \equiv b \pmod{W}}} v_y e(\alpha(xy - b)/W) \sum_{\substack{d|xy+2 \\ d \leq D}} w_d \right| \right)^2 \\
& \leq \left(\sum_{X \leq x \leq X'} |u_x|^2 \right) \left(\sum_{X \leq x \leq X'} \left| \sum_{\substack{Y \leq y \leq Y' \\ Z \leq xy \leq Z' \\ xy \equiv b \pmod{W}}} v_y e(\alpha(xy - b)/W) \sum_{\substack{d|xy+2 \\ d \leq D}} w_d \right|^2 \right) \\
& \ll X(\log X)^{6^A} \sum_{\substack{1 \leq b' \leq W, (b', W)=1 \\ X \leq x \leq X' \\ x \equiv b' \pmod{W} \\ d_1, d_2 \leq D}} |w_{d_1}| |w_{d_2}| \prod_{i=1}^2 \left| \sum_{\substack{Y \leq y_i \leq Y' \\ Z \leq xy_i \leq Z' \\ y_i b' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i}}} v_{y_i} e(\alpha xy_i/W) \right|.
\end{aligned}$$

By the partial summation,

$$\begin{aligned}
& \sum_{\substack{Y \leq y_i \leq Y' \\ Z \leq xy_i \leq Z' \\ y_i b' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i}}} v_{y_i} e(\alpha xy_i/W) \\
& = v_{Y'} \sum_{\substack{1 \leq y_i \leq Y' \\ Z \leq xy_i \leq Z' \\ y_i b' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i}}} e(\alpha xy_i/W) - v_Y \sum_{\substack{1 \leq y_i \leq Y-1 \\ Z \leq xy_i \leq Z' \\ y_i b' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i}}} e(\alpha xy_i/W) \\
& \quad - \sum_{Y \leq Y'' \leq Y'-1} (v_{Y''+1} - v_{Y''}) \sum_{\substack{1 \leq y_i \leq Y'' \\ Z \leq xy_i \leq Z' \\ y_i b' \equiv b \pmod{W} \\ xy_i \equiv -2 \pmod{d_i}}} e(\alpha xy_i/W) \\
& \ll \left(v_{Y'} + v_Y + \sum_{Y \leq Y'' \leq Y'-1} (v_{Y''+1} - v_{Y''}) \right) \min \left\{ \frac{Y}{[d_i, W]}, \frac{1}{\|\alpha[d_i, W]x/W\|} \right\} \\
& \ll_A (\log(XY))^A \min \left\{ \frac{Y}{[d_i, W]}, \frac{1}{\|\alpha[d_i, W]x/W\|} \right\}.
\end{aligned}$$

For any $1 \leq b' \leq W$ with $(b', W) = 1$,

$$\begin{aligned}
& \sum_{\substack{X \leq x \leq X' \\ x \equiv b' \pmod{W} \\ d_1, d_2 \leq D}} |w_{d_1}| |w_{d_2}| \prod_{i=1}^2 \min \left\{ \frac{Y}{[d_i, W]}, \frac{1}{\|\alpha[d_i, W]x/W\|} \right\} \\
& \leq (\log(XY))^{2A} \sum_{\substack{X \leq x \leq X' \\ x \equiv b' \pmod{W}}} \left(\sum_{d \leq D} \tau(d)^A \min \left\{ \frac{Y}{[d, W]}, \frac{1}{\|\alpha[d, W]x/W\|} \right\} \right)^2 \\
& \leq (\log(XY))^{2A} \sum_{\substack{X \leq x \leq X' \\ x \equiv b' \pmod{W}}} \left(\sum_{h \leq D} \tau(hW)^{A+1} \min \left\{ \frac{Y}{hW}, \frac{1}{\|\alpha hx\|} \right\} \right)^2 \\
& \quad (\text{where } h = [d, W]/W) \\
& \leq (\log(XY))^{2A} \sum_{\substack{X \leq x \leq X' \\ x \equiv b' \pmod{W}}} \left(\sum_{2^k \leq D} \sum_{2^k \leq h \leq 2^{k+1}} \tau(hW)^{A+1} \min \left\{ \frac{Y}{hW}, \frac{1}{\|\alpha hx\|} \right\} \right)^2 \\
& \ll (\log(DXY))^{2A+2} \max_{H \leq D} \sum_{\substack{X \leq x \leq X' \\ x \equiv b' \pmod{W}}} \left(\sum_{H/2 \leq h \leq H} \tau(hW)^{A+1} \min \left\{ \frac{Y}{HW}, \frac{1}{\|\alpha hx\|} \right\} \right)^2.
\end{aligned}$$

And for any $H \leq D$,

$$\begin{aligned}
& \sum_{\substack{X \leq x \leq X' \\ x \equiv b' \pmod{W}}} \left(\sum_{H/2 \leq h \leq H} \tau(hW)^{A+1} \min \left\{ \frac{Y}{HW}, \frac{1}{\|\alpha hx\|} \right\} \right)^2 \\
& \leq \sum_{\substack{X \leq x \leq X' \\ x \equiv b' \pmod{W}}} \left(\tau(W)^{2A+2} \sum_{h \leq H} \tau(h)^{2A+2} \right) \left(\frac{Y}{HW} \sum_{h \leq H} \min \left\{ \frac{Y}{HW}, \frac{1}{\|\alpha hx\|} \right\} \right) \\
& \ll_A (\log D)^{2^{2A+2}} Y \sum_{1 \leq x \leq X'H} \tau(x) \min \left\{ \frac{Y}{HW}, \frac{1}{\|\alpha x\|} \right\}.
\end{aligned}$$

Finally,

$$\begin{aligned}
& \sum_{1 \leq x \leq X'H} \tau(x) \min \left\{ \frac{Y}{HW}, \frac{1}{\|\alpha x\|} \right\} \\
& \ll \left(\sum_{1 \leq x \leq X'H} \tau(x)^2 \right)^{1/2} \left(\frac{Y}{HW} \sum_{1 \leq x \leq X'H} \min \left\{ \frac{X'Y}{xW}, \frac{1}{\|\alpha x\|} \right\} \right)^{1/2} \\
& \ll \left(\frac{X^2 Y^2 (\log(DXq))^4}{W^2} \left(\frac{1}{q} + \frac{DW}{Y} + \frac{qW}{XY} \right) \right)^{1/2}.
\end{aligned}$$

□

Define

$$\tau_k(x) = \{(d_1, d_2, \dots, d_k) : d_1 d_2 \cdots d_k \mid x\}.$$

Let $G(x)$ be an arbitrary complex function over $\mathbb{N}$. Consider

Type I sums

$$\sum_{\substack{M < m \leq M_1 \\ L < l \leq L_1 \\ P \leq ml \leq P_1}} a_m G(ml) \quad \text{and} \quad \sum_{\substack{M < m \leq M_1 \\ L < l \leq L_1 \\ P \leq ml \leq P_1}} a_m (\log l) G(ml)$$

where $M_1 \leq 2M$, $L_1 \leq 2L$, $|a_m| \leq \tau_5(m) \log P$,

and

Type II sums

$$\sum_{\substack{M < m \leq M_1 \\ L < l \leq L_1 \\ P \leq ml \leq P_1}} a_m b_l G(ml)$$

where $M_1 \leq 2M$, $L_1 \leq 2L$, $|a_m| \leq \tau_5(m) \log P$, $|b_l| \leq \tau_5(l) \log P$.

The following Lemma is due to Heath-Brown [7]:

Lemma 2.4. *Let P, P_1, u, v, z be positive integers satisfying $2 < P < P_1 \leq 2P$, $2 \leq u < v \leq z \leq P$, $u^2 \leq z$, $128uz^2 \leq P_1$, $2^{18}P_1 \leq v^3$. Then we may decompose the sum*

$$\sum_{P < n \leq P_1} \Lambda(n) G(n)$$

into $O((\log P)^6)$ sums, each of which is either of type I with $L \geq z$ or of type II with $u \leq L \leq v$.

Lemma 2.5. *For any $\alpha \in \mathfrak{m}$,*

$$S_{n, z_0}^\pm(\alpha) \ll n(\log n)^{68-B/4}.$$

Proof. Clearly

$$S_{n, z_0}^\pm(\alpha) = \sum_{\substack{n^{0.99} \leq p \leq n \\ p \equiv b \pmod{W}}} e(\alpha(p-b)/W) \log p \sum_{d|(p+2, P(z_0))} \lambda_D^\pm(d) + O(n^{0.995}).$$

And notice that for any $x \leq n$,

$$\left| \sum_{d|(x+2, P(z_0))} \lambda_D^\pm(d) \right| \leq \tau(x+2) \ll_\epsilon n^\epsilon.$$

So it suffices to estimate the sum

$$\sum_{\substack{n' \leq x \leq n \\ x \equiv b \pmod{W}}} \Lambda(x) e(\alpha(x-b)/W) \sum_{d|(x+2, P(z_0))} \lambda_D^\pm(d), \quad (2.6)$$

where $n' \geq n/2$. Since $\alpha \in \mathfrak{m}$, there exist $1 \leq a \leq q$ with $(a, q) = 1$ and $(\log n)^B \leq q \leq n(\log n)^{-B}$ such that $|\alpha q - a| \leq (\log n)^B/n$. Applying Lemma 2.4

with $u = n^{0.17}$, $v = n^{0.34}$ and $z = n^{0.35}$, the sum (2.6) can be decomposed into $O((\log n)^6)$ type I sums

$$\sum_{\substack{M < m \leq M_1 \\ L < l \leq L_1 \\ ml \equiv b \pmod{W} \\ n' \leq ml \leq n}} a_m e(\alpha(ml - b)/W) \sum_{d|(ml+2, P(z_0))} \lambda_D^\pm(d)$$

and

$$\sum_{\substack{M < m \leq M_1 \\ L < l \leq L_1 \\ ml \equiv b \pmod{W} \\ n' \leq ml \leq n}} a_m (\log l) e(\alpha(ml - b)/W) \sum_{d|(ml+2, P(z_0))} \lambda_D^\pm(d)$$

with $L \geq n^{0.35}$, and type II sums

$$\sum_{\substack{M < m \leq M_1 \\ L < l \leq L_1 \\ ml \equiv b \pmod{W} \\ n' \leq ml \leq n}} a_m b_l e(\alpha(ml - b)/W) \sum_{d|(ml+2, P(z_0))} \lambda_D^\pm(d)$$

with $n^{0.17} \leq L \leq n^{0.34}$. Noting that $\lambda_D^\pm(d) = 0$ whenever $d \geq D$, in view of (2.4) with $A = 5$, these type II sums are all $\ll n(\log n)^{2^{13}-B/4}$. And by (2.5), all type I sums are $\ll n(\log n)^{67-B/4}$.

□

3. THE MAJOR ARCS

Define

$$\Delta(x; q) := \max_{\substack{1 \leq r \leq q \\ (r, q) = 1}} \left| \sum_{\substack{p \leq x \\ p \equiv r \pmod{q}}} \log p - \frac{x}{\phi(q)} \right|.$$

The well-known Bombieri-Vinogradov theorem asserts that for any $A > 0$

$$\sum_{q \leq n^{1/2-\epsilon}} \max_{x \leq n} \Delta(x; q) \ll_{A, \epsilon} \frac{n}{(\log n)^A}. \quad (3.1)$$

Define

$$\phi_2(q) := q \prod_{2 < p|q} \left(1 - \frac{2}{p}\right).$$

Lemma 3.1.

$$S_{n, z_0}^+(0) \leq \frac{4e^{-\gamma} k_0 \mathfrak{S}_1 n}{\phi_2(W) \log n} (F(s) + O(e^{-s} (\log n)^{-1/3}))$$

and

$$S_{n, z_0}^-(0) \geq \frac{4e^{-\gamma} k_0 \mathfrak{S}_1 n}{\phi_2(W) \log n} (f(s) + O(e^{-s} (\log n)^{-1/3})),$$

where γ is Euler's constant,

$$\mathfrak{S}_1 = \prod_{p > 2} \left(1 - \frac{1}{(p-1)^2}\right) = 0.6601 \dots$$

and $s = \log D / \log z_0$.

Proof.

$$\begin{aligned} S_{n,z_0}^+(0) &= \sum_{\substack{d|P(z_0) \\ b \equiv -2 \pmod{(d,W)}}} \lambda_D^+(d) \sum_{\substack{p \leq n \\ p \equiv b \pmod{W} \\ p \equiv -2 \pmod{d}}} \log p \\ &= \sum_{\substack{d|P(z_0) \\ (d,W)=1}} \lambda_D^+(d) \left(\frac{n}{\phi(Wd)} + O(\Delta(n; Wd)) \right) \end{aligned}$$

since $(W, b+2) = 1$. Since $\lambda_D^+(d)$ vanishes for $d \geq D$, by (3.1) we have,

$$S_{n,z_0}^+(0) = \frac{n}{\phi(W)} \sum_{\substack{d|P(z_0) \\ (d,W)=1}} \frac{\lambda_D^+(d)}{\phi(d)} + O\left(\frac{n}{(\log n)^{5B}}\right).$$

Applying (2.1),

$$\sum_{\substack{d|P(z_0) \\ (d,W)=1}} \frac{\lambda_D^+(d)}{\phi(d)} \leq \prod_{p \leq z_0, p \nmid W} \left(1 - \frac{1}{p-1}\right) (F(s) + O(e^{-s}(\log D)^{-1/3})).$$

Similarly,

$$S_{n,z_0}^-(0) \geq \frac{n}{\phi(W)} \prod_{p \leq z_0, p \nmid W} \left(1 - \frac{1}{p-1}\right) (f(s) + O(e^{-s}(\log n)^{-1/3})).$$

Finally, by the Mertens theorem,

$$\prod_{p \leq z_0, p \nmid W} \left(1 - \frac{1}{p-1}\right)^{-1} = \frac{\phi_2(W)}{2\phi(W)} \prod_{2 < p \leq z_0} \left(1 - \frac{1}{p-1}\right)^{-1} = \frac{\phi_2(W)}{4\phi(W)} (\mathfrak{S}_1^{-1} e^\gamma \log z_0 + O(1)).$$

□

Let $m = (n-b)/W$. Define $\Lambda_*(x) = \log x$ or 0 according to whether x is prime.

Lemma 3.2. *Suppose that $1 \leq a \leq q \leq (\log n)^B$ and $(a, q) = 1$. Then we have*

$$\left| S_{n,z_0}(a/q) - \frac{\mathbf{1}_{(W,q)=1} \mu(q) \tau^*(a, q) 4e^{-\gamma} k_0 \mathfrak{S}_1 W m}{\phi_2(Wq) \log(Wm+b)} \right| \leq \frac{5e^{-\gamma} k_0 \mathfrak{S}_1 (F(s) - f(s)) W m}{\phi_2(W) \log(Wm+b)}, \quad (3.2)$$

where $\mathbf{1}_{(W,q)=1} = 1$ or 0 according to whether $(W, q) = 1$,

$$\tau^*(a, q) = \sum_{\substack{d|q \\ (d,q/d)=1}} e(ar_d/q),$$

and $1 \leq r_d \leq q$ is the unique integer r such that $Wr \equiv -b \pmod{d}$ and $Wr \equiv -b-2 \pmod{q/d}$.

Proof. Clearly

$$\sum_{\substack{1 \leq x \leq m \\ (Wx+b+2, P(z_0))=1}} \Lambda_*(Wx+b)e(ax/q) = \sum_{\substack{1 \leq r \leq q \\ (Wr+b, q)=1 \\ (Wr+b+2, q)=1}} e(ar/q) \sum_{\substack{1 \leq p \leq Wm+b \\ p \equiv Wr+b \pmod{Wq} \\ (p+2, P(z_0))=1}} \log p + O(\log^{B+1} n).$$

Notice that

$$\begin{aligned} & \sum_{\substack{1 \leq p \leq Wm+b \\ p \equiv Wr+b \pmod{Wq}}} \log p \sum_{d|(p+2, P(z_0))} \lambda_D^-(d) \\ & \leq \sum_{\substack{1 \leq p \leq Wm+b \\ p \equiv Wr+b \pmod{Wq}}} \log p \sum_{d|(p+2, P(z_0))} \mu(d) \\ & \leq \sum_{\substack{1 \leq p \leq Wm+b \\ p \equiv Wr+b \pmod{Wq}}} \log p \sum_{d|(p+2, P(z_0))} \lambda_D^+(d). \end{aligned}$$

From the proof of Lemma 3.1, we know that

$$\begin{aligned} & \left| \sum_{\substack{1 \leq p \leq Wm+b \\ p \equiv Wr+b \pmod{Wq} \\ (p+2, P(z_0))=1}} \log p - \frac{e^{-\gamma} k_0 \mathfrak{S}_1 Wm}{\phi_2(Wq) \log(Wm+b)} \right| \\ & \leq \frac{1.1 e^{-\gamma} k_0 \mathfrak{S}_1 (F(s) - f(s)) Wm}{\phi_2(Wq) \log(Wm+b)}. \end{aligned}$$

By noting that W is even and $(W, b(b+2)) = 1$,

$$\begin{aligned} \sum_{\substack{1 \leq r \leq q \\ (Wr+b, q)=1 \\ (Wr+b+2, q)=1}} e(ar/q) &= \sum_{d_1, d_2 | q} \mu(d_1) \mu(d_2) \sum_{\substack{1 \leq r \leq q \\ d_1 | Wr+b \\ d_2 | Wr+b+2}} e(ar/q) \\ &= \sum_{\substack{d_1 d_2 = q \\ (d_1, d_2)=1 \\ (d_1, W)=(d_2, W)=1}} \mu(d_1) \mu(d_2) e(ar_{d_1}/q) \\ &= \begin{cases} \mu(q) \tau^*(a, q) & \text{if } (W, q) = 1, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Furthermore, we have

$$|\{1 \leq r \leq q : ((Wr+b)(Wr+b+2), q) = 1\}| = q \prod_{p|q, p \nmid W} \left(1 - \frac{2}{p}\right) = \frac{\phi_2(Wq)}{\phi_2(W)}.$$

All are done. $\square$

Lemma 3.3. Suppose that $1 \leq a \leq q \leq (\log n)^B$ and $(a, q) = 1$. Then for any $\alpha \in \mathfrak{M}_{a, q}$,

$$S_{n, z_0}^\pm(\alpha) = \frac{S_{n, z_0}^\pm(a/q)}{m} \sum_{1 \leq y \leq m} e(\theta y) + O\left(\frac{m}{(\log m)^{3B}}\right), \quad (3.3)$$

where $\theta = \alpha - a/q$.

Proof. By the partial summation,

$$\begin{aligned} & S_{n,z_0}^{\pm}(\alpha) \\ &= e(\theta m) \sum_{1 \leq x \leq m} \Lambda_*(Wx + b)e(ax/q) \sum_{d|(Wx+b+2, P(z_0))} \lambda_D^{\pm}(d) \\ & \quad - \sum_{y \leq m-1} (e(\theta(y+1)) - e(\theta y)) \sum_{1 \leq x \leq y} \Lambda_*(Wx + b)e(ax/q) \sum_{d|(Wx+b+2, P(z_0))} \lambda_D^{\pm}(d). \end{aligned}$$

Recalling that $(W, b+2) = 1$, write

$$\begin{aligned} & \sum_{1 \leq x \leq y} \Lambda_*(Wx + b)e(ax/q) \sum_{d|(Wx+b+2, P(z_0))} \lambda_D^{\pm}(d) \\ &= \sum_{\substack{d|P(z_0) \\ (d,W)=1 \\ d \leq D}} \lambda_D^{\pm}(d) \sum_{\substack{1 \leq x \leq y \\ Wx \equiv -b-2 \pmod{d}}} \Lambda_*(Wx + b)e(ax/q) \\ &= \sum_{\substack{d|P(z_0) \\ (d,W)=1 \\ d \leq D}} \lambda_D^{\pm}(d) \left(\sum_{\substack{1 \leq r \leq q \\ (Wr+b, q)=1 \\ Wr \equiv -b-2 \pmod{(d,q)}}} e(ar/q) \sum_{\substack{1 \leq x \leq y \\ x \equiv r \pmod{q} \\ Wx \equiv -b-2 \pmod{d}}} \Lambda_*(Wx + b) + O(\log^{B+1} n) \right). \end{aligned}$$

Now

$$\sum_{\substack{1 \leq x \leq y \\ x \equiv r \pmod{q} \\ Wx \equiv -b-2 \pmod{d}}} \Lambda_*(Wx + b) = \frac{Wy + b}{\phi(W[d, q])} + O(\Delta(Wy + b; W[d, q])).$$

Notice that for any d' with $q \mid d'$, $|\{d : [d, q] = d'\}| \leq \tau(q)$. Hence

$$\begin{aligned} & \sum_{1 \leq x \leq y} \Lambda_*(Wx + b)e(ax/q) \sum_{d|(Wx+b+2, P(z_0))} \lambda_D^{\pm}(d) \\ &= \sum_{\substack{d|P(z_0) \\ (d,W)=1 \\ d \leq D}} \lambda_D^{\pm}(d) \sum_{\substack{1 \leq r \leq q \\ (Wr+b, q)=1 \\ Wr \equiv -b-2 \pmod{(d,q)}}} e(ar/q) \frac{Wy + b}{\phi(W[d, q])} + O\left(\frac{Wy + b}{(\log(Wy + b))^{5B}}\right). \end{aligned}$$

So

$$S_{n,z_0}^{\pm}(\alpha) = W \sum_{1 \leq y \leq m} e(\theta y) \sum_{\substack{d|P(z_0) \\ (d,W)=1 \\ d \leq D}} \lambda_D^{\pm}(d) \sum_{\substack{1 \leq r \leq q \\ (Wr+b, q)=1 \\ Wr \equiv -b-2 \pmod{(d,q)}}} \frac{e(ar/q)}{\phi(W[d, q])} + O\left(\frac{m}{(\log m)^{3B}}\right).$$

Setting $\theta \rightarrow 0$ in the above equation, we obtain that

$$W \sum_{\substack{d|P(z_0) \\ (d,W)=1 \\ d \leq D}} \lambda_D^{\pm}(d) \sum_{\substack{1 \leq r \leq q \\ (Wr+b, q)=1 \\ Wr \equiv -b-2 \pmod{(d,q)}}} \frac{e(ar/q)}{\phi(W[d, q])} = \frac{S_{n,z_0}^{\pm}(a/q)}{m} + O\left(\frac{1}{(\log m)^{3B}}\right).$$

□

Combining Lemmas 2.2, 2.5, 3.1, 3.2 and 3.3, we get

Lemma 3.4. *Suppose that $1 \leq a \leq q \leq (\log n)^B$ and $(a, q) = 1$. Then for any $\alpha \in \mathfrak{M}_{a,q}$,*

$$\left| S_{n,z_0}(\alpha) - \frac{\mathbf{1}_{(W,q)=1} \mu(q) \tau^*(a, q) 4e^{-\gamma} k_0 \mathfrak{S}_1 W}{\phi_2(Wq) \log n} \sum_{1 \leq y \leq m} e(\theta y) \right| \leq \frac{15e^{-\gamma} k_0 \mathfrak{S}_1 (F(s) - f(s)) n}{\phi_2(W) \log n}, \quad (3.4)$$

where $\theta = \alpha - a/q$. Furthermore, for any $\alpha \in \mathfrak{m}$,

$$|S_{n,z_0}(\alpha)| \leq \frac{5e^{-\gamma} k_0 \mathfrak{S}_1 (F(s) - f(s)) n}{\phi_2(W) \log n}. \quad (3.5)$$

Lemma 3.5.

$$\sum_{\substack{p_1, p_2 \leq n \\ (p_i+2, P(z_0))=1 \\ p_i \equiv b \pmod{W} \\ p_2 - p_1 = WM}} 1 \ll \frac{k_0^2 n W}{\phi_2(W)^2 \log^4 n} \prod_{\substack{p|M \\ p \nmid W}} \left(1 + \frac{2}{p}\right) \prod_{\substack{p|(WM+2)(WM-2) \\ p \nmid W}} \left(1 + \frac{1}{p}\right). \quad (3.6)$$

Proof. Let $z_1 = n^{1/10}$. Let ω_1 and ω_2 be two multiplicative functions satisfying that

$$\omega_1(p) = \begin{cases} 4 & \text{if } p < z_0 \text{ and } p \nmid WM(WM-2)(WM+2), \\ 3 & \text{if } p < z_0 \text{ and } p \mid (WM-2)(WM+2), \ p \nmid W, \\ 2 & \text{if } p < z_0, \ p \mid M \text{ and } p \nmid W, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$\omega_2(p) = \begin{cases} 2 & \text{if } z_0 \leq p < z_1 \text{ and } p \nmid WM, \\ 1 & \text{if } z_0 \leq p < z_1, \ p \mid M \text{ and } p \nmid W, \\ 0 & \text{otherwise,} \end{cases}$$

for prime p . And for $1 \leq i \leq 2$, let g_i be the multiplicative functions with

$$g_i(p) = \frac{\omega_i(p)}{p} \left(1 - \frac{\omega_i(p)}{p}\right)^{-1}$$

for prime p , and let

$$G_1^{(i)}(z) = \sum_{\substack{l|P(z) \\ l < z}} g_i(l).$$

Define

$$\lambda_1(d) = \frac{d}{\omega_1(d)} \sum_{\substack{l|P(z_0) \\ d|l < z_0}} \frac{\mu(l/d) \mu(l) g_1(l)}{G_1^{(1)}(z_0)}$$

for $d \mid P(z_0)$ and

$$\lambda_2(d) = \frac{d}{\omega_2(d)} \sum_{\substack{l|P(z_0, z_1) \\ d|l < z_1}} \frac{\mu(l/d) \mu(l) g_2(l)}{G_1^{(2)}(z_1)}$$

for $d \mid P(z_0, z_1) = \prod_{z_0 \leq p < z_1} p$. Then $\lambda_1(1) = \lambda_2(1) = 1$. Therefore

$$\begin{aligned}
& \sum_{\substack{x_1, x_2 \leq n/W \\ (Wx_i + b + 2, P(z_0)) = 1 \\ (Wx_i + b, P(z_1)) = 1 \\ x_2 - x_1 = M}} 1 \\
& \leq \sum_{x \leq n/W} \left(\sum_{\substack{d \mid P(z_0) \\ d \mid (Wx+b)(Wx+WM+b) \\ d \mid (Wx+b+2)(Wx+WM+b+2)}} \lambda_1(d) \right)^2 \left(\sum_{\substack{d \mid P(z_0, z_1) \\ d \mid (Wx+b)(Wx+WM+b)}} \lambda_2(d) \right)^2 \\
& = \sum_{\substack{d_1, d_2 \mid P(z_0) \\ d_3, d_4 \mid P(z_0, z_1)}} \lambda_1(d_1) \lambda_1(d_2) \lambda_2(d_3) \lambda_2(d_4) \sum_{\substack{x \leq n/W \\ [d_3, d_4] \mid (Wx+b)(Wx+WM+b) \\ [d_1, d_2], [d_3, d_4] \mid (Wx+b+2)(Wx+WM+b+2)}} 1 \\
& = \sum_{\substack{d_1, d_2 \mid P(z_0) \\ d_3, d_4 \mid P(z_0, z_1)}} \lambda_1(d_1) \lambda_1(d_2) \lambda_2(d_3) \lambda_2(d_4) \omega_1([d_1, d_2]) \omega_2([d_3, d_4]) \left(\frac{n/W}{[d_1, d_2][d_3, d_4]} + O(1) \right).
\end{aligned}$$

By Selberg's sieve method, we know that $|\lambda_1(d)|, |\lambda_2(d)| \leq 1$ and

$$\begin{aligned}
& \sum_{\substack{d_1, d_2 \mid P(z_0) \\ d_3, d_4 \mid P(z_0, z_1)}} \lambda_1(d_1) \lambda_1(d_2) \lambda_2(d_3) \lambda_2(d_4) \frac{\omega_1([d_1, d_2]) \omega_2([d_3, d_4])}{[d_1, d_2][d_3, d_4]} \\
& = \left(\sum_{d_1, d_2 \mid P(z_0)} \lambda_1(d_1) \lambda_1(d_2) \frac{\omega_1([d_1, d_2])}{[d_1, d_2]} \right) \left(\sum_{d_3, d_4 \mid P(z_0, z_1)} \lambda_2(d_3) \lambda_2(d_4) \frac{\omega_2([d_3, d_4])}{[d_3, d_4]} \right) \\
& = \frac{1}{G_1^{(1)}(z_0)} \frac{1}{G_1^{(2)}(z_1)} \ll \prod_{p < z_0} \left(1 - \frac{\omega_1(p)}{p} \right) \prod_{z_0 \leq p < z_1} \left(1 - \frac{\omega_2(p)}{p} \right).
\end{aligned}$$

Thus

$$\begin{aligned}
& \sum_{\substack{x_1, x_2 \leq n/W \\ (Wx_i + b + 2, P(z_0)) = 1 \\ (Wx_i + b, P(z_1)) = 1 \\ x_2 - x_1 = M}} 1 \\
& \ll \frac{n}{W} \prod_{p \mid P(z_0)} \left(1 - \frac{\omega_1(p)}{p} \right) \prod_{p \mid P(z_0, z_1)} \left(1 - \frac{\omega_2(p)}{p} \right) \\
& \ll \frac{n}{W(\log z_0)^2 (\log z_1)^2} \prod_{p \mid W} \left(1 + \frac{4}{p} \right) \prod_{\substack{p \mid M \\ p \nmid W}} \left(1 + \frac{2}{p} \right) \prod_{\substack{p \mid (WM+2)(WM-2) \\ p \nmid W}} \left(1 + \frac{1}{p} \right) \\
& \ll \frac{k_0^2 n W}{\phi_2(W)^2 \log^4 n} \prod_{\substack{p \mid M \\ p \nmid W}} \left(1 + \frac{2}{p} \right) \prod_{\substack{p \mid (WM+2)(WM-2) \\ p \nmid W}} \left(1 + \frac{1}{p} \right).
\end{aligned}$$

□

Lemma 3.6. *Then*

$$\sum_{\substack{p_1, p_2, p_3, p_4 \leq n \\ (p_i + 2, P(z_0)) = 1 \\ p_i \equiv b \pmod{W} \\ p_1 + p_4 = p_2 + p_3}} 1 \ll \frac{k_0^4 W n^3}{\phi_2(W)^4 \log^8 n}. \quad (3.7)$$

Proof. Applying Lemma 3.5,

$$\begin{aligned} \sum_{\substack{p_1, p_2, p_3, p_4 \leq n \\ (p_i + 2, P(z_0)) = 1 \\ p_i \equiv b \pmod{W} \\ p_1 + p_4 = p_2 + p_3}} 1 &\leq \sum_{2 < M \leq n/W} \left(\sum_{\substack{p_1, p_2 \leq n \\ (p_i + 2, P(z_0)) = 1 \\ p_i \equiv b \pmod{W} \\ p_1 + WM = p_2}} 1 \right)^2 + O(n^2) \\ &\ll \frac{k_0^4 n^2 W^2}{\phi_2(W)^4 \log^8 n} \sum_{2 < M \leq n/W} \prod_{\substack{p|M \\ p \nmid W}} \left(1 + \frac{2}{p}\right)^2 \prod_{\substack{p|(WM-2)(WM+2) \\ p \nmid W}} \left(1 + \frac{1}{p}\right)^2. \end{aligned}$$

By the Hölder inequality,

$$\begin{aligned} &\sum_{2 < M \leq n/W} \prod_{\substack{p|M \\ p \nmid W}} \left(1 + \frac{2}{p}\right)^2 \prod_{\substack{p|(WM-2)(WM+2) \\ p \nmid W}} \left(1 + \frac{1}{p}\right)^2 \\ &\leq \left(\sum_{2 < M \leq n/W} \prod_{p|M} \left(1 + \frac{2}{p}\right)^6 \right)^{1/3} \prod_{j=1}^2 \left(\sum_{2 < M \leq n/W} \prod_{\substack{p|WM+2(-1)^j \\ p \neq 2}} \left(1 + \frac{1}{p}\right)^6 \right)^{1/3}. \end{aligned}$$

Since $(1 + p^{-1})^6 \leq 1 + 24p^{-1}$,

$$\begin{aligned} \sum_{2 < M \leq n/W} \prod_{\substack{p|WM \pm 2 \\ p \neq 2}} \left(1 + \frac{1}{p}\right)^6 &\leq \sum_{2 < M \leq n/W} \sum_{\substack{d|WM \pm 2 \\ 2 \nmid d}} \frac{\tau(d)^{12}}{d} \\ &= \sum_{\substack{d|n \pm 2 \\ (d, W) = 1}} \frac{\tau(d)^{12}}{d} \sum_{\substack{2 < M \leq n/W \\ d|WM \pm 2}} 1 \ll \frac{n}{W}. \end{aligned}$$

And by [3, Lemma 14], we have

$$\sum_{2 < M \leq n/W} \prod_{p|M} \left(1 + \frac{2}{p}\right)^6 \leq \sum_{2 < M \leq n/W} \prod_{p|M} \left(1 + \frac{1}{p}\right)^{12} \ll \frac{n}{W}.$$

□

4. PROOF OF THEOREM 1.1

First, let us introduce Green and Tao's enveloping sieve. Let N be a large integer. Suppose that $a_1, \dots, a_k, b_1, \dots, b_k$ be integers with $|a_i|, |b_i| \leq N$. We say

$$\mathcal{F}(x) := \prod_{i=1}^k (a_i x + b_i).$$

is a k -linear form.

For every integer $q \geq 1$, define

$$\gamma_{\mathcal{F}}(p) := q^{-1} |\{1 \leq x \leq q : (\mathcal{F}(x), q) = 1\}| > 0.$$

Let

$$X_{R!}(x) = \{x \in \mathbb{Z} : (\mathcal{F}(x), R!) = 1\},$$

where $1 \leq R \leq N$.

Lemma 4.1 ([5, Proposition 3.1]). *There exists a non-negative function $\beta_R : \mathbb{Z} \rightarrow \mathbb{R}$ satisfying the following properties:*

(i)

$$\beta_R(x) \gg_k \mathfrak{S}_{\mathcal{F}}^{-1} \log^k R \mathbf{1}_{X_{R!}}(x) \quad (4.1)$$

for all integers n , where

$$\mathfrak{S}_{\mathcal{F}} := \prod_p \frac{\gamma_{\mathcal{F}}(p)}{(1 - 1/p)^k}.$$

(ii)

$$\beta_R(x) \ll_{k,\epsilon} N^{\epsilon} \quad (4.2)$$

for all $1 \leq x \leq N$ and $\epsilon > 0$.

(iii)

$$\beta_R(x) = \sum_{q \leq R^2} \sum_{\substack{1 \leq a \leq q \\ (a,q)=1}} w(a/q) e(-ax/q), \quad (4.3)$$

where $w(a/q) = w_R(a/q)$ satisfies $w(1) = 1$ and

$$|w(a/q)| \ll_{k,\epsilon} q^{\epsilon-1}$$

for all $1 \leq a \leq q \leq R^2$ with $(a, q) = 1$.

(iv) For $1 \leq a \leq q \leq R^2$ with $(a, q) = 1$, if q is not square-free, or $\gamma(q) = 1$ and $q > 1$, then $w(a/q) = 0$.

Green and Tao also established a restriction theorem for β_R :

Lemma 4.2 ([5, Proposition 4.2]). *Let R, N be large numbers such that $1 \leq R \leq N^{1/10}$. Let $k, \mathcal{F}, \beta_R$ as defined in Lemma 4.1. Suppose that $\{u_i\}_{i=1}^N$ is an arbitrary sequence of complex numbers. Then for any $\rho > 2$,*

$$\left(\sum_{r \in \mathbb{Z}_N} \left| \frac{1}{N} \sum_{1 \leq x \leq N} u_x \beta_R(x) e(-xr/N) \right|^\rho \right)^{1/\rho} \ll_{\rho,k} \left(\frac{1}{N} \sum_{1 \leq x \leq N} |u_x|^2 \beta_R(x) \right)^{1/2}. \quad (4.4)$$

The following lemma can be derived by a trivial modification of Chen's original proof in [1]:

Lemma 4.3. *Suppose that W is a positive integer. Then there exists a function $n_0(W)$ such that for every $1 \leq b \leq W$ with $(b(b+2), W) = 1$ and $n \geq n_0(W)$,*

$$|\{p \leq n : p \equiv b \pmod{W}, p+2 \in \mathcal{P}_2 \text{ and } (p+2, P(n^{1/10})) = 1\}| \geq \frac{C_1}{\phi_2(W)} \frac{n}{(\log n)^2},$$

where C_1 is an absolute constant.

Lemma 4.4. *Suppose that $n_0(W)$ is an increasing positive function for all positive integers W , and $G(n)$ is an increasing positive function with $\lim_{n \rightarrow \infty} G(n) = \infty$. Then there exists an increasing positive function $W_0(n)$ for sufficiently large n such that $n \geq n_0(W(n))$, $W_0(n) \leq G(n)$ and $\lim_{n \rightarrow \infty} W_0(n) = \infty$.*

Proof. Let

$$W_0(n) = \max\{W \leq G(n) : n_0(W) \leq n\}.$$

Clearly $W_0(n)$ is well-defined for sufficiently large n . Assume on the contrary that $\lim_{n \rightarrow \infty} W_0(n) < \infty$, i.e., there exists an integer W' such that $W_0(n) < W'$ for all n . Let $n' \geq n_0(W')$ be an integer such that $G(n') \geq W'$. Obviously such n' exists. But $W_0(n') \geq W'$ now. This leads to a contradiction. $\square$

Let C_2 be the implied constant in (4.1) with $k = 2$. Let

$$\varpi = \frac{\min\{C_1 C_2, 1\}}{10000}.$$

Let C_3 be the implied constant in (4.4) with $\rho = 12/5$ and $k = 2$. And let C_4 be the implied constant in (3.7).

Lemma 4.5. *There exists $0 < \kappa \leq \varpi$ such that we can choose $0 < \delta, \epsilon \leq 1$ with $\epsilon^{6C_3^{12/5}\delta^{-12/5}+60C_4\delta^{-4}} \geq \kappa$ and*

$$3072\epsilon^2(C_3\delta^{-12/5} + 5C_4\delta^{-4}) + 72C_3^{24/13}C_4^{3/13}\delta^{1/13} \leq \varpi^6.$$

Proof. Let

$$\delta = \min\left\{\frac{\varpi^{78}}{144^{13}C_3^{24}C_4^3}, 1\right\}$$

and

$$\epsilon = \min\left\{\frac{\varpi^3\delta^2}{192(C_3\delta^{-12/5} + 5C_4\delta^{-4})^{1/2}}, 1\right\}.$$

Clearly $3072\epsilon^2(C_3\delta^{-12/5} + 5C_4\delta^{-4}) + 72C_3^{24/13}C_4^{3/13}\delta^{1/13} \leq \varpi^6$. So we may arbitrarily choose a κ satisfying

$$\kappa \leq \min\{\epsilon^{6C_3^{12/5}\delta^{-12/5}+60C_4\delta^{-4}}, \varpi\}.$$

$\square$

Let κ be a small constant satisfying the requirements of Lemma 4.5. Notice that $f(s)$ is increasing, $F(s)$ is decreasing and $F(s), f(s) = 1 + O(e^{-s})$. Choose a sufficiently large k_0 satisfying that

$$20(F(k_0/4) - f(k_0/4)) \leq \kappa^2.$$

Suppose that n is a sufficiently large integer. Let $w = w(n)$ be a positive function satisfying $P(w) \leq \log n$, $n \geq n_0(P(w))$ (where n_0 is defined in Lemma 4.3) and $\lim_{n \rightarrow \infty} w(n) = \infty$. By Lemma 4.4, such w exists. Let $W = P(w)$. The following lemma can be easily verified:

Lemma 4.6. *For any odd integer n with $3 \mid n$, there exist $1 \leq b_1, b_2, b_3 \leq W$ with $(b_i(b_i + 2), W) = 1$ such that*

$$n \equiv b_1 + b_2 + b_3 \pmod{W}.$$

Furthermore, if $n \equiv 4 \pmod{6}$, then there exist $1 \leq b_1, b_2 \leq W$ with $(b_i(b_i + 2), W) = 1$ such that

$$n \equiv b_1 + b_2 \pmod{W}.$$

Now suppose that n is odd. Suppose that $1 \leq b_1, b_2, b_3 \leq W$ are integers satisfying $(b_i(b_i + 2), W) = 1$ and $n \equiv b_1 + b_2 + b_3 \pmod{W}$. Let $n' = (n - b_1 - b_2 - b_3)/W$.

Let N be a prime in the interval $[(1 + \kappa^2/20)n/W, (1 + \kappa^2/10)n/W]$ and $R = N^{1/10}$. Thanks to the prime number theorem, such a prime N always exists whenever n is sufficiently large. Let $\mathcal{F}_i(x) = (Wx + b_i)(Wx + b_i + 2)$ for $i = 1, 2$. Substituting $N, R, \mathcal{F}_i$ to Lemma 4.1, we get the desired functions $\beta_i = \beta_{i,R}$ for $i = 1, 2$.

Let

$A_i = \{x \leq (n - b_i)/2W : Wx + b_i \text{ is Chen's prime and } (Wx + b_i + 2, P(n^{1/10})) = 1\}$, for $i = 1, 2$, and define

$$\mathfrak{a}_i(x) = \mathbf{1}_{A_i}(x) \frac{C_2 \mathfrak{S}_1^{-1} \phi_2(W) (\log(Wx + b_i))^2}{1000 n}.$$

By Lemma 4.3, clearly we have

$$\sum_x \mathfrak{a}_i(x) \geq (1 - \kappa) \frac{C_2 \mathfrak{S}_1^{-1} \phi_2(W) (\log n)^2 |A_i|}{1000 n} \geq 6\varpi, \quad (4.5)$$

whenever n is sufficiently large.

On the other hand, it is easy to see that

$$\mathfrak{S}_{\mathcal{F}_i} = \prod_{\substack{p|W \\ p>2}} \frac{1}{(1 - 1/p)^2} \prod_{p \nmid W} \frac{1 - 2/p}{(1 - 1/p)^2} = \prod_{\substack{p|W \\ p>2}} \frac{p}{p-2} \prod_{p>2} \left(1 - \frac{1}{(p-1)^2}\right) = \frac{W \mathfrak{S}_1}{\phi_2(W)}.$$

Hence by (4.1), for any x

$$\beta_i(x) \geq \frac{C_2 \phi_2(W) \mathfrak{S}_1^{-1}}{W} (\log R)^2 \mathbf{1}_{A_i}(x) \geq N \mathfrak{a}_i(x). \quad (4.6)$$

Let

$$A_3 = \{x \leq (n - b_3)/W : Wx + b_3 \text{ is prime and } (Wx + b_3 + 2, P(n^{1/k_0})) = 1\}.$$

and define

$$\mathfrak{a}_3(x) = \mathbf{1}_{A_3}(x) \frac{e^\gamma \phi_2(W) \log(Wx + b_3) \log n}{4k_0 \mathfrak{S}_1 n}.$$

In view of Lemmas 2.2 and 3.1, we also have

$$\sum_x \mathfrak{a}_3(x) = \frac{e^\gamma \phi_2(W) \log n}{4k_0 \mathfrak{S}_1 n} S_{n, n^{1/k_0}}(0) \in [1 - \kappa^2, 1 + \kappa^2]. \quad (4.7)$$

Below we identify the set $\{1, 2, \dots, N\}$ with the group $\mathbb{Z}_N = \mathbb{Z}/N\mathbb{Z}$. If there exist $x_1 \in A_1$, $x_2 \in A_2$ and $x_3 \in A_3$ satisfying $x_1 + x_2 + x_3 = n'$ in $\mathbb{Z}_N$, then the equality also holds in $\mathbb{Z}$. In fact, since $x_1 + x_2 \leq n/W$ and $x_3 < n/W$ in $\mathbb{Z}$, we must have $x_1 + x_2 + x_3 < n' + N$ in $\mathbb{Z}$.

For any function $f : \mathbb{Z}_N \rightarrow \mathbb{C}$, define

$$\tilde{f}(r) = \sum_{x \in \mathbb{Z}_N} f(x) e(-xr/N).$$

Lemma 4.7. *Let $\nu_i = \beta_i/N$, then for any $r \in \mathbb{Z}_N$,*

$$|\tilde{\nu}_i(r) - \mathbf{1}_{r=0}| \leq C_5 w^{-1/2},$$

where C_5 is an absolute constant and $\mathbf{1}_{r=0} = 1$ or 0 according to whether $r = 0$.

Proof. See [5, Lemma 6.1]. (Notice that our definitions are a little different from Green and Tao's.) $\square$

Lemma 4.8. *For any $0 \neq r \in \mathbb{Z}_N$,*

$$|\tilde{\mathbf{a}}_3(r)| \leq \frac{2}{w-2} + 0.9\kappa^2.$$

Proof. If $r/N \in \mathfrak{m}$, then by Lemma 3.4, we have

$$|\tilde{\mathbf{a}}_3(r)| = \frac{e^\gamma \phi_2(W) \log n}{k_0 \mathfrak{S}_1 n} |S_{n, n^{1/k_0}}(r/N)| \leq \frac{2}{5} \kappa^2.$$

Suppose that there exist $1 \leq a \leq q \leq (\log n)^B$ with $(a, q) = 1$ such that $r/N \in \mathfrak{M}_{a,q}$. Then applying Lemma 3.4,

$$|\tilde{\mathbf{a}}_3(r)| \leq \frac{\phi_2(W) \log n}{k_0 \mathfrak{S}_1 n} \left(\frac{\mathbf{1}_{(W,q)=1} |\tau^*(a, q)| k_0 \mathfrak{S}_1 W}{\phi_2(Wq) \log n} \left| \sum_{1 \leq y \leq m} e(\theta y) \right| + \frac{7k_0 \mathfrak{S}_1 \kappa^2 n}{10\phi_2(W) \log n} \right),$$

where $m = (n - b)/W$ and $\theta = r/N - a/q$. Recall that $\tau^*(a, q) = 0$ whenever $(W, q) > 1$. And if $a = q = 1$, since $r \neq 0$,

$$\left| \sum_{1 \leq y \leq m} e(yr/N) \right| \leq \left| \sum_{1 \leq y \leq N} e(yr/N) \right| + N - m \leq \frac{\kappa^2}{10} N.$$

Suppose that $q > 1$ and $(W, q) = 1$. Then by noting $W = \prod_{p < w} p$,

$$\frac{\phi_2(W) |\tau^*(a, q)|}{\phi_2(Wq)} \leq \frac{\phi_2(W) |\{d \mid q : (d, q/d) = 1\}|}{\phi_2(Wq)} \leq \frac{2}{w-2},$$

since q has at least one prime divisor not less than w . Finally, we have $WN \leq 1.1n$. $\square$

Suppose that $\delta, \epsilon > 0$ are two small numbers to be chosen later. For $1 \leq i \leq 3$, let

$$\mathcal{R}_i = \{r \in \mathbb{Z}_N : |\tilde{a}_i(r)| > \delta\},$$

$$\mathcal{B}_i = \{x \in \mathbb{Z}_N : \|xr/N\| \leq \epsilon\},$$

and define $\mathfrak{b}_i = \mathbf{1}_{\mathcal{B}_i}/|B_i|$.

Lemma 4.9.

$$|\mathcal{R}_3| \leq 60C_4\delta^{-4}.$$

Proof. By (3.7),

$$\begin{aligned} \sum_{r \in \mathbb{Z}_N} |\tilde{\mathbf{a}}_3(r)|^4 &= N \sum_{\substack{x_1, x_2, x_3, x_4 \in \mathbb{Z}_N \\ x_1 + x_4 = x_2 + x_3}} \mathbf{a}_3(x_1) \mathbf{a}_3(x_2) \mathbf{a}_3(x_3) \mathbf{a}_3(x_4) \\ &\leq N \left(\frac{e^\gamma \phi_2(W) (\log n)^2}{4k_0 \mathfrak{S}_1 n} \right)^4 \sum_{\substack{p_1, p_2, p_3, p_4 \leq n \\ (p_i + 2, P(n^{1/k_0})) = 1 \\ p_i \equiv b_3 \pmod{W} \\ p_1 + p_4 = p_2 + p_3}} 1 \\ &\leq C_4 e^{4\gamma} (4\mathfrak{S}_1)^{-4} \frac{NW}{n}. \end{aligned}$$

Hence

$$|\mathcal{R}_3| \leq \sum_{r \in \mathbb{Z}_N} \delta^{-4} |\tilde{\mathbf{a}}_3(r)|^4 \leq e^{4\gamma} (4\mathfrak{S}_1)^{-4}.$$

□

Lemma 4.10.

$$|\mathcal{R}_1|, |\mathcal{R}_2| \leq 6C_3^{12/5} \delta^{-12/5}$$

provided that $w \geq C_5^2$.

Proof. Suppose that $i \in \{1, 2\}$. Let $u_x = \mathbf{a}_i(x)/\nu_i(x)$ or 0 according to whether $\nu_i(x) \neq 0$. By (4.6), clearly $0 \leq u_x \leq 1$. If $w \geq C_5^2$, by Lemma 4.7 we have $\tilde{\nu}_i(0) \leq 2$. Applying (4.4),

$$\begin{aligned} \sum_{r \in \mathbb{Z}_N} \left| \sum_{1 \leq x \leq N} u_x \nu_i(x) e(-xr/N) \right|^{12/5} &\leq C_3^{12/5} \left(\sum_{1 \leq x \leq N} |u_x|^2 \nu_i(x) \right)^{6/5} \\ &\leq C_3^{12/5} |\tilde{\nu}_i(0)|^{6/5} \\ &\leq 6C_3^{12/5}. \end{aligned}$$

It follows that $|\mathcal{R}_i| \leq 6C_3^{12/5} \delta^{-12/5}$.

□

Lemma 4.11.

$$|\mathcal{B}_i| \geq e^{|\mathcal{R}_i|} N.$$

Proof. This is a simple application of the pigeonhole principle (cf. [14, Lemma 1.4]).

□

For two functions $f, g : \mathbb{Z}_N \rightarrow \mathbb{C}$. Define

$$f * g(x) = \sum_{y \in \mathbb{Z}_N} f(y) g(x - y).$$

It is easy to check that $\widetilde{(f * g)} = \tilde{f} \tilde{g}$. Let $\mathbf{a}'_i = \mathbf{a}_i * \mathbf{b}_i * \mathbf{b}_i$ for $1 \leq i \leq 3$.

Lemma 4.12. *Suppose that $\epsilon^{|\mathcal{R}_3|} \geq (2/(w-2) + 0.9\kappa^2)\kappa^{-1}$. Then for any $x \in \mathbb{Z}_N$,*

$$|\mathbf{a}'_3(x)| \leq \frac{1+2\kappa}{N}.$$

Proof.

$$\begin{aligned} |\mathbf{a}'_3(x)| &= |\mathbf{a}_3(x) * \mathbf{b}_3(x) * \mathbf{b}_3(x)| \\ &\leq \frac{1}{N} |\tilde{\mathbf{a}}_3(0)| |\tilde{\mathbf{b}}_3(0)|^2 + \frac{1}{N} \sum_{r \neq 0} |\tilde{\mathbf{a}}_3(r) \tilde{\mathbf{b}}_3(r)^2| \\ &\leq \frac{1}{N} |\tilde{\mathbf{a}}_3(0)| |\tilde{\mathbf{b}}_3(0)|^2 + \frac{1}{N} \max_{r \neq 0} |\tilde{\mathbf{a}}_3(r)| \sum_{r \in \mathbb{Z}_N} |\tilde{\mathbf{b}}_3(r)|^2 \\ &\leq \frac{1+\kappa^2}{N} + \frac{1}{|B_3|} \left(\frac{2}{w-2} + 0.9\kappa^2 \right), \end{aligned}$$

where we used Lemma 4.8 in the last step. Thus the desired result easily follows from Lemma 4.11. $\square$

Lemma 4.13. *Let $i \in \{1, 2\}$. Suppose that $\epsilon^{|\mathcal{R}_i|} \geq C_5 \kappa^{-1} w^{-1/2}$ and $C_5 w^{-1/2} \leq \kappa$. Then for any $x \in \mathbb{Z}_N$,*

$$|\mathbf{a}'_i(x)| \leq \frac{1+2\kappa}{N}.$$

Proof. Since $\nu_i(x) \geq \mathbf{a}_i(x)$, applying Lemma 4.7

$$\begin{aligned} |\mathbf{a}'_i(x)| &\leq |\nu_i(x) * \mathbf{b}_i(x) * \mathbf{b}_i(x)| \\ &\leq \frac{1}{N} |\tilde{\nu}_i(0)| |\tilde{\mathbf{b}}_i(0)|^2 + \frac{1}{N} \max_{r \neq 0} |\tilde{\nu}_i(r)| \sum_{r \in \mathbb{Z}_N} |\tilde{\mathbf{b}}_i(r)|^2 \\ &\leq \frac{1 + C_5 w^{-1/2}}{N} + \frac{C_5 w^{-1/2}}{|B_i|}. \end{aligned}$$

We are done. $\square$

Lemma 4.14. *For $1 \leq i \leq 3$,*

$$|1 - \tilde{\mathbf{b}}_i(r)| \leq 16\epsilon^2$$

for any $r \in \mathcal{R}_i$.

Proof. See the proof of Lemma 6.7 of [4]. $\square$

Lemma 4.15. *Suppose that $w \geq C_5^2$. Then*

$$\begin{aligned} &\left| \sum_{\substack{x_1, x_2, x_3 \in \mathbb{Z}_N \\ x_1 + x_2 + x_3 = n'}} \mathbf{a}'_1(x_1) \mathbf{a}'_2(x_2) \mathbf{a}'_3(x_3) - \sum_{\substack{x_1, x_2, x_3 \in \mathbb{Z}_N \\ x_1 + x_2 + x_3 = n'}} \mathbf{a}_1(x_1) \mathbf{a}_2(x_2) \mathbf{a}_3(x_3) \right| \\ &\leq \frac{3072\epsilon^2 (C_3^{12/5} \delta^{-12/5} + 5C_4 \delta^{-4}) + 72C_3^{24/13} C_4^{3/13} \delta^{1/13}}{N}. \end{aligned}$$

Proof. Clearly

$$\sum_{\substack{x_1, x_2, x_3 \in \mathbb{Z}_N \\ x_1 + x_2 + x_3 = n'}} \mathbf{a}_1(x_1) \mathbf{a}_2(x_2) \mathbf{a}_3(x_3) = \frac{1}{N} \sum_{r \in \mathbb{Z}_N} \tilde{\mathbf{a}}_1(r) \tilde{\mathbf{a}}_2(r) \tilde{\mathbf{a}}_3(r) e(n'r/N).$$

Hence

$$\begin{aligned} & \left| \sum_{\substack{x_1, x_2, x_3 \in \mathbb{Z}_N \\ x_1 + x_2 + x_3 = n'}} \mathbf{a}'_1(x_1) \mathbf{a}'_2(x_2) \mathbf{a}'_3(x_3) - \sum_{\substack{x_1, x_2, x_3 \in \mathbb{Z}_N \\ x_1 + x_2 + x_3 = n'}} \mathbf{a}_1(x_1) \mathbf{a}_2(x_2) \mathbf{a}_3(x_3) \right| \\ & \leq \frac{1}{N} \sum_{r \in \mathbb{Z}_N} |\tilde{\mathbf{a}}_1(r) \tilde{\mathbf{a}}_2(r) \tilde{\mathbf{a}}_3(r) (1 - \tilde{\mathbf{b}}_1(r)^2 \tilde{\mathbf{b}}_2(r)^2 \tilde{\mathbf{b}}_3(r)^2)|. \end{aligned}$$

Since $w \geq C_5^2$, for $i = 1, 2$,

$$|\tilde{a}_i(r)| \leq |\tilde{a}_i(0)| \leq |\tilde{\nu}_i(0)| \leq 2.$$

We also have

$$|\tilde{a}_3(r)| \leq |\tilde{a}_3(0)| \leq 1 + \kappa^2.$$

If $r \in \mathcal{R}_1 \cap \mathcal{R}_2 \cap \mathcal{R}_3$, then by Lemma 4.14,

$$|1 - \tilde{\mathbf{b}}_i(r)^2| \leq |1 - \tilde{\mathbf{b}}_i(r)| (1 + |\tilde{\mathbf{b}}_i(r)|) \leq |1 - \tilde{\mathbf{b}}_i(r)| (1 + |\tilde{\mathbf{b}}_i(0)|) \leq 32\epsilon^2.$$

So

$$\begin{aligned} & |1 - \tilde{\mathbf{b}}_1(r)^2 \tilde{\mathbf{b}}_2(r)^2 \tilde{\mathbf{b}}_3(r)^2| \\ & \leq |1 - \tilde{\mathbf{b}}_1(r)^2| + |\tilde{\mathbf{b}}_1(r)|^2 |1 - \tilde{\mathbf{b}}_2(r)^2| + |\tilde{\mathbf{b}}_1(r) \tilde{\mathbf{b}}_2(r)|^2 |1 - \tilde{\mathbf{b}}_3(r)^2| \\ & \leq 96\epsilon^2. \end{aligned}$$

Thus

$$\begin{aligned} \sum_{r \in \mathcal{R}_1 \cap \mathcal{R}_2 \cap \mathcal{R}_3} |\tilde{\mathbf{a}}_1(r) \tilde{\mathbf{a}}_2(r) \tilde{\mathbf{a}}_3(r) (1 - \tilde{\mathbf{b}}_1(r)^2 \tilde{\mathbf{b}}_2(r)^2 \tilde{\mathbf{b}}_3(r)^2)| & \leq 768\epsilon^2 \min_{1 \leq i \leq 3} |\mathcal{R}_i| \\ & \leq 256\epsilon^2 (12C_3^{12/5} \delta^{-12/5} + 60C_4 \delta^{-4}). \end{aligned}$$

On the other hand, by the Hölder inequality,

$$\begin{aligned} & \sum_{r \notin \mathcal{R}_1 \cap \mathcal{R}_2 \cap \mathcal{R}_3} |\tilde{\mathbf{a}}_1(r) \tilde{\mathbf{a}}_2(r) \tilde{\mathbf{a}}_3(r) (1 - \tilde{\mathbf{b}}_1(r)^2 \tilde{\mathbf{b}}_2(r)^2 \tilde{\mathbf{b}}_3(r)^2)| \\ & \leq 2 \sum_{r \notin \mathcal{R}_1 \cap \mathcal{R}_2 \cap \mathcal{R}_3} |\tilde{\mathbf{a}}_1(r) \tilde{\mathbf{a}}_2(r) \tilde{\mathbf{a}}_3(r)| \\ & \leq 2 \max_{r \notin \mathcal{R}_1 \cap \mathcal{R}_2 \cap \mathcal{R}_3} |\tilde{\mathbf{a}}_1(r) \tilde{\mathbf{a}}_2(r) \tilde{\mathbf{a}}_3(r)|^{1/13} \sum_{r \in \mathbb{Z}_N} |\tilde{\mathbf{a}}_1(r) \tilde{\mathbf{a}}_2(r) \tilde{\mathbf{a}}_3(r)|^{12/13} \\ & \leq 4\delta^{1/13} \left(\sum_{r \in \mathbb{Z}_N} |\tilde{\mathbf{a}}_1(r)|^{12/5} \right)^{5/13} \left(\sum_{r \in \mathbb{Z}_N} |\tilde{\mathbf{a}}_2(r)|^{12/5} \right)^{5/13} \left(\sum_{r \in \mathbb{Z}_N} |\tilde{\mathbf{a}}_3(r)|^4 \right)^{3/13} \\ & \leq 4\delta^{1/13} (6C_3^{12/5})^{10/13} (60C_4)^{3/13}. \end{aligned}$$

□

Lemma 4.16. *Suppose that $0 < \theta_1, \theta_2, \theta_3 \leq 1$ with $\theta_1 + \theta_2 + \theta_3 > 1$. Let*

$$\theta = \min\{\theta_1, \theta_2, \theta_3, (\theta_1 + \theta_2 + \theta_3 - 1)/4\}.$$

Suppose that N is a prime greater than $2\theta^{-2}$, and X_1, X_2, X_3 are subsets of $\mathbb{Z}_N$ with $|X_i| \geq \theta_i N$. Then for any $y \in \mathbb{Z}_N$, we have

$$|\{(x_1, x_2, x_3) : x_i \in X_i, x_1 + x_2 + x_3 = y\}| \geq \theta^3 N^2.$$

Proof. See [11, Lemma 3.3]. □

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. For $1 \leq i \leq 3$, let

$$A'_i = \{x \in \mathbb{Z}_N : \mathfrak{a}'_i(x) \geq \varpi/N\}.$$

In view of (4.7) and Lemma 4.12,

$$\frac{1+2\kappa}{N}|A'_3| + \frac{\varpi}{N}(N - |A'_3|) \geq \sum_{x \in \mathbb{Z}_N} \mathfrak{a}'_3(x) = \sum_{x \in \mathbb{Z}_N} \mathfrak{a}_3(x) \geq 1 - \kappa^2.$$

Therefore $|A'_3| \geq (1 - 3\varpi)N$. Similarly, by (4.5) and Lemma 4.13, for $i = 1, 2$

$$\frac{1+2\kappa}{N}|A'_i| + \frac{\varpi}{N}(N - |A'_i|) \geq \sum_{x \in \mathbb{Z}_N} \mathfrak{a}_i(x) \geq 6\varpi.$$

Hence $|A'_1|, |A'_2| \geq \frac{9}{2}\varpi N$. Since $|A'_1| + |A'_2| + |A'_3| \geq (1 + 6\varpi)N$, with the help of Lemma 4.16,

$$\sum_{\substack{x_1 \in A'_1, x_2 \in A'_2, x_3 \in A'_3 \\ x_1 + x_2 + x_3 = n'}} 1 \geq 2\varpi^3 N^2.$$

By Lemma 4.5, we may choose $0 < \delta, \epsilon \leq 1$ with $\epsilon^{6C_3^{12/5}\delta^{-12/5} + 60C_4\delta^{-4}} \geq \kappa$ satisfying

$$3072\epsilon^2(C_3^{12/5}\delta^{-12/5} + 5C_4\delta^{-4}) + 72C_3^{24/13}C_4^{3/13}\delta^{1/13} \leq \varpi^6.$$

Notice that $w = w(n)$ tends to infinity with n . So we may assume that $w \geq \max\{20\kappa^{-2} + 2, C_5^2\kappa^{-4}\}$. Since $\max_{1 \leq i \leq 3} |\mathcal{R}_i| \leq 6C_3^{12/5}\delta^{-12/5} + 6C_4\delta^{-4}$, the requirements of Lemmas 4.12, 4.13 and 4.15 are obviously satisfied.

Applying Lemma 4.15,

$$\begin{aligned} & \sum_{\substack{x_1, x_2, x_3 \in \mathbb{Z}_N \\ x_1 + x_2 + x_3 = n'}} \mathfrak{a}_1(x_1)\mathfrak{a}_2(x_2)\mathfrak{a}_3(x_3) \\ & \geq \sum_{\substack{x_1, x_2, x_3 \in \mathbb{Z}_N \\ x_1 + x_2 + x_3 = n'}} \mathfrak{a}'_1(x_1)\mathfrak{a}'_2(x_2)\mathfrak{a}'_3(x_3) - \frac{3072\epsilon^2(C_3^{12/5}\delta^{-12/5} + 5C_4\delta^{-4}) + 72C_3^{24/13}C_4^{3/13}\delta^{1/13}}{N} \\ & \geq \sum_{\substack{x_1 \in A'_1, x_2 \in A'_2, x_3 \in A'_3 \\ x_1 + x_2 + x_3 = n'}} \left(\frac{\varpi}{N}\right)^3 - \frac{\varpi^6}{N} \\ & \geq \frac{\varpi^6}{N}. \end{aligned}$$

This completes the proof. □

REFERENCES

- [1] J.-R. Chen, *On the representation of a large even integer as the sum of a prime and a product of at most two primes*, Sci. Sinica, **16**(1973), 157-176.
- [2] J. G. van der Corput, *Über Summen von Primzahlen und Primzahlquadraten*, Math. Ann., **116**(1939), 1-50.
- [3] B. Green, *On arithmetic structures in dense sets of integers*, Duke Math. J., **114**(2002), 215-238.
- [4] B. Green, *Roth's theorem in the primes*, Ann. Math., **161**(2005), 1609-1636.
- [5] B. Green and T. Tao, *Restriction theory of the Selberg sieve, with application*, J. Theor. Nombres Bordeaux, **18** (2006), 147-182.
- [6] B. Green and T. Tao, *The primes contain arbitrarily long arithmetic progressions*, Ann. Math., **167**(2008), 481-547.
- [7] D. R. Heath-Brown, *Prime numbers in short intervals and a generalized Vaughan identity*, Canad. J. Math., **35** (1982), 1365-1377.
- [8] H. Iwaniec, *Rosser's sieve*, Acta Arith., **36** (1980), 171-202.
- [9] H. Iwaniec, *A new form of the error term in the linear sieve*, Acta Arith., **37** (1980), 307-320.
- [10] H. Iwaniec, *Sieve methods*, Graduate course, Rutgers, 1996.
- [11] H.-Z. Li and H. Pan, *Ternary Goldbach problem for the subsets of primes with positive relative densities*, preprint, arXiv:math.NT/0701240.
- [12] X. Meng, *A mean value theorem on the binary Goldbach problem and its application*, Monatsh. Math., **151** (2007), 319-332.
- [13] T. P. Peneva, *On the ternary Goldbach problem with primes p_i such that $p_i + 2$ are almost-primes*, Acta Math. Hungar., **86** (2000), 305-318.
- [14] T. Tao, *The Roth-Bourgain Theorem*, preprint, unpublished.
- [15] D. I. Tolev, *Arithmetic progressions of prime-almost-prime twins*, Acta Arith., **88** (1999), 67-98.
- [16] D. I. Tolev, *Additive problems with prime numbers of special type*, Acta Arith., **96** (2000), 53-88.
- [17] R. C. Vaughan, *The Hardy-Littlewood Method*, Second edition, Cambridge University Press, Cambridge, 1997.
- [18] I. M. Vinogradov, *The representation of an odd number as a sum of three primes*, Dokl. Akad. Nauk. SSSR., **16**(1937), 139-142.

DEPARTMENT OF MATHEMATICS, SHANGHAI JIAOTONG UNIVERSITY, SHANGHAI 200240,
PEOPLE'S REPUBLIC OF CHINA

E-mail address: lih@sjtu.edu.cn

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S RE-
PUBLIC OF CHINA

E-mail address: haopan79@yahoo.com.cn